

## Product datasheet for RC217417L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MASP1 (NM\_001031849) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MASP1 (NM\_001031849) Human Tagged ORF Clone Lentiviral Particle

Symbol: MASP1

Synonyms: 3MC1; CRARF; CRARF1; MAP-1; MAP1; MAP4; MASP; MASP-3; MASP3; PRSS5; RaRF

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001031849

ORF Size: 1140 bp

**ORF Nucleotide** 

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC217417).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001031849.1</u>

 RefSeq Size:
 2072 bp

 RefSeq ORF:
 1143 bp

 Locus ID:
 5648

 UniProt ID:
 P48740

 Cytogenetics:
 3q27.3

**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Complement and coagulation cascades





ORIGENE

MW: 43.5 kDa

**Gene Summary:** 

This gene encodes a serine protease that functions as a component of the lectin pathway of complement activation. The complement pathway plays an essential role in the innate and adaptive immune response. The encoded protein is synthesized as a zymogen and is activated when it complexes with the pathogen recognition molecules of lectin pathway, the mannose-binding lectin and the ficolins. This protein is not directly involved in complement activation but may play a role as an amplifier of complement activation by cleaving complement C2 or by activating another complement serine protease, MASP-2. The encoded protein is also able to cleave fibrinogen and factor XIII and may may be involved in coagulation. A splice variant of this gene which lacks the serine protease domain functions as an inhibitor of the complement pathway. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Apr 2010]