

## Product datasheet for RC217503L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ST2 (IL1RL1) (NM\_016232) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: ST2 (IL1RL1) (NM 016232) Human Tagged ORF Clone Lentiviral Particle

Symbol: ST2

Synonyms: DER4; FIT-1; IL33R; ST2; ST2L; ST2V; T1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_016232 **ORF Size:** 1668 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC217503).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 016232.4

 RefSeq Size:
 2058 bp

 RefSeq ORF:
 1671 bp

 Locus ID:
 9173

 UniProt ID:
 Q01638

 Cytogenetics:
 2q12.1

Domains: TIR, ig, IGc2, IG

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane





ORÏGENE

**MW:** 63.36 kDa

**Gene Summary:** The protein encoded by this gene is a member of the interleukin 1 receptor family. Studies of

the similar gene in mouse suggested that this receptor can be induced by proinflammatory stimuli, and may be involved in the function of helper T cells. This gene, interleukin 1 receptor, type I (IL1R1), interleukin 1 receptor, type II (IL1R2) and interleukin 1 receptor-like 2 (IL1RL2) form a cytokine receptor gene cluster in a region mapped to chromosome 2q12. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul

2008]