

Product datasheet for RC217522L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TRIF (TICAM1) (NM_182919) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TRIF (TICAM1) (NM_182919) Human Tagged ORF Clone Lentiviral Particle

Symbol: TRIF

Synonyms: IIAE6; MyD88-3; PRVTIRB; TICAM-1; TRIF

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM_182919

ORF Size: 2136 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC217522).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 182919.1

 RefSeq Size:
 2460 bp

 RefSeq ORF:
 2139 bp

 Locus ID:
 148022


 UniProt ID:
 Q8IUC6

 Cytogenetics:
 19p13.3

Protein Families: Druggable Genome

Protein Pathways: Toll-like receptor signaling pathway





ORIGENE

MW: 76.2 kDa

Gene Summary:

This gene encodes an adaptor protein containing a Toll/interleukin-1 receptor (TIR) homology domain, which is an intracellular signaling domain that mediates protein-protein interactions between the Toll-like receptors (TLRs) and signal-transduction components. This protein is involved in native immunity against invading pathogens. It specifically interacts with toll-like receptor 3, but not with other TLRs, and this association mediates dsRNA induction of interferon-beta through activation of nuclear factor kappa-B, during an antiviral immune response. Mutations in this gene are associated with encephalopathy, acute, infection-induced. [provided by RefSeq, Jul 2020]