

## Product datasheet for RC217551L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NDST3 (NM\_004784) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** NDST3 (NM\_004784) Human Tagged ORF Clone Lentiviral Particle

Symbol: NDST3
Synonyms: HSST3

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_004784 **ORF Size:** 2619 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC217551).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 004784.1

 RefSeq Size:
 3188 bp

 RefSeq ORF:
 2622 bp

 Locus ID:
 9348

 UniProt ID:
 095803

 Cytogenetics:
 4q26

**Domains:** Sulfotransfer

**Protein Families:** Transmembrane





## NDST3 (NM\_004784) Human Tagged ORF Clone Lentiviral Particle - RC217551L2V

**Protein Pathways:** Heparan sulfate biosynthesis, Metabolic pathways

MW: 100.7 kDa

**Gene Summary:** This gene encodes a member of the heparan sulfate/heparin GlcNAc N-deacetylase/ N-

sulfotransferase family. The encoded enzyme is a type II transmembrane protein that resides in the Golgi apparatus. This monomeric bifunctional enzyme catalyzes the N-deacetylation and N-sulfation of N-acetylglucosamine residues in heparan sulfate and heparin, which are

the initial chemical modifications required for the biosynthesis of the functional

oligosaccharide sequences that define the specific ligand binding activities of heparan sulfate

and heparin. [provided by RefSeq, Nov 2008]