

## Product datasheet for RC217577L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rad51L1 (RAD51B) (NM\_002877) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Rad51L1 (RAD51B) (NM\_002877) Human Tagged ORF Clone Lentiviral Particle

Symbol: Rad51L1

Synonyms: R51H2; RAD51L1; REC2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_002877 **ORF Size:** 1050 bp

**ORF Nucleotide** 

OTI Disclaimer:

.030 56

Sequence:

The ORF insert of this clone is exactly the same as(RC217577).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002877.4

 RefSeq Size:
 1875 bp

 RefSeq ORF:
 1053 bp

 Locus ID:
 5890

 UniProt ID:
 015315

 Cytogenetics:
 14q24.1

Domains: AAA

**Protein Families:** Druggable Genome





**Protein Pathways:** Homologous recombination

MW: 38.1 kDa

**Gene Summary:** The protein encoded by this gene is a member of the RAD51 protein family. RAD51 family

members are evolutionarily conserved proteins essential for DNA repair by homologous recombination. This protein has been shown to form a stable heterodimer with the family member RAD51C, which further interacts with the other family members, such as RAD51, XRCC2, and XRCC3. Overexpression of this gene was found to cause cell cycle G1 delay and

cell apoptosis, which suggested a role of this protein in sensing DNA damage.

Rearrangements between this locus and high mobility group AT-hook 2 (HMGA2, GeneID 8091) have been observed in uterine leiomyomata. [provided by RefSeq, Mar 2016]