

## Product datasheet for RC217592L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Caspase 10 (CASP10) (NM 001230) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Caspase 10 (CASP10) (NM\_001230) Human Tagged ORF Clone Lentiviral Particle

Symbol: CASP10

Synonyms: ALPS2; FLICE-2; FLICE2; MCH4

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_001230 **ORF Size:** 1437 bp

**ORF Nucleotide** 

1 137 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC217592).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001230.3

RefSeq Size: 3651 bp
RefSeq ORF: 1440 bp

Locus ID: 843

 UniProt ID:
 Q92851

 Cytogenetics:
 2q33.1

**Domains:** Peptidase\_C14, DED, CASc

**Protein Families:** Druggable Genome, Protease





**Protein Pathways:** Apoptosis, RIG-I-like receptor signaling pathway

**MW:** 54.4 kDa

**Gene Summary:** This gene encodes a protein which is a member of the cysteine-aspartic acid protease

(caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and

the protein itself is processed by caspase 8. Mutations in this gene are associated with type IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Apr 2011]