

Product datasheet for RC217608L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Germinal Center Kinase (MAP4K2) (NM_004579) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Germinal Center Kinase (MAP4K2) (NM_004579) Human Tagged ORF Clone Lentiviral Particle

Symbol: Germinal Center Kinase

Synonyms: BL44; GCK; RAB8IP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_004579

ORF Size: 2460 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC217608).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004579.2

 RefSeq Size:
 2964 bp

 RefSeq ORF:
 2463 bp

 Locus ID:
 5871

 UniProt ID:
 Q12851

Cytogenetics: 11q13.1

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: MAPK signaling pathway





Germinal Center Kinase (MAP4K2) (NM_004579) Human Tagged ORF Clone Lentiviral Particle – RC217608L3V

MW:

91.4 kDa

Gene Summary:

The protein encoded by this gene is a member of the serine/threonine protein kinase family. Although this kinase is found in many tissues, its expression in lymphoid follicles is restricted to the cells of germinal centre, where it may participate in B-cell differentiation. This kinase can be activated by TNF-alpha, and has been shown to specifically activate MAP kinases. This kinase is also found to interact with TNF receptor-associated factor 2 (TRAF2), which is involved in the activation of MAP3K1/MEKK1. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015]