

Product datasheet for RC217620L4

SLC5A8 (NM_145913) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Tag: mGFP

Symbol: SLC5A8

Synonyms: AIT; SMCT; SMCT1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as (RC217620).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:

Cloning sites used for ORF Shuttling:



The diagram illustrates the pET28a(+) vector structure. It features a series of restriction sites: EcoRI, BamHI, RBS, SgfI, MluI, NotI, and XhoI. The mGFP Tag sequence is inserted between the SgfI and MluI sites. The insertion site is highlighted with a green bar, and the mGFP Tag sequence is also shown in green. The vector ends with an ORF (Open Reading Frame) sequence.

----- NNN ACG CGT ACG CGG CCG CTC GAG ATG AGC GGG GGC -----
 T R T R P L E M S G G - - -
 ----- GGA CTC AGA GTT TGG GTA GGA AGC
 C G A T T G A A T T G

* The last cedar before the Stop cedar of the QRF.

ACCN: NM 145913

ORF Size: 1830 bp




[View online »](#)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2025 OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, USA

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_145913.2
RefSeq Size:	3286 bp
RefSeq ORF:	1833 bp
Locus ID:	160728
UniProt ID:	Q8N695
Cytogenetics:	12q23.1-q23.2
Domains:	SSF
Protein Families:	Transmembrane
MW:	66.6 kDa
Gene Summary:	SLC5A8 has been shown to transport iodide by a passive mechanism (Rodriguez et al., 2002 [PubMed 12107270]) and monocarboxylates and short-chain fatty acids by a sodium-coupled mechanism (Gopal et al., 2004 [PubMed 15322102]). In kidney, SLC5A8 functions as a high-affinity sodium-coupled lactate transporter involved in reabsorption of lactate and maintenance of blood lactate levels (Thangaraju et al., 2006 [PubMed 16873376]).[supplied by OMIM, Dec 2008]

Product images:

Circular map for RC217620L4