

## Product datasheet for RC217665

### H2AZ2 (NM\_138635) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** H2AZ2 (NM\_138635) Human Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** H2AZ2  
**Synonyms:** H2A.Z-2; H2AFV; H2AV  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**ORF Nucleotide Sequence:** >RC217665 representing NM\_138635  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCTGGAGGCAAAGCTGGAAGGACAGTGGGAAGGCCAAGGCTAAGGCAGTATCTCGCTCACAGAGAG  
CTGGGCTACAGTTTCTGTGGCCGCATCCACAGACACTTGAAGACTCGCACCACAAGCCATGGAAGGGT  
GGGTGCCACTGCTGCCGTGTACAGTGTGCGATTCTGGAGTACCTCACTGCAGAGGTGCTGGAGCTGGCA  
GGTAATGCTTCTAAGGATCTCAAAGTAAAGCGTATCACTCCGCGTCACTGCAGCTTGAATCCGTGGT  
ATGAAGAGTTGGATTCTTATCAAGGCTACCATAGCTGGGGTGAGAAGAGAAGGTGTTCT

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC217665 representing NM\_138635  
Red=Cloning site Green=Tags(s)  
MAGGKAGKDSGKAKAKAVSRSQRAGLQFPVGRHRLKTRTTSHGRVGATAAVYSAAILEYLTAEVLELA  
GNASKDLKVKRITPRHLQLAIRGDEELDSLKATIAGGEKRRCS

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:** [https://cdn.origene.com/chromatograms/mk6481\\_h01.zip](https://cdn.origene.com/chromatograms/mk6481_h01.zip)

**Restriction Sites:** SgfI-MluI



[View online »](#)

**Cloning Scheme:**


**ACCN:** NM\_138635

**ORF Size:** 342 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [NM\\_138635.3](#)

**RefSeq Size:** 825 bp

**RefSeq ORF:** 345 bp

Locus ID: 94239

UniProt ID: [Q71UI9](#)

Cytogenetics: 7p13

Domains: histone


Protein Families: Druggable Genome

Protein Pathways: Systemic lupus erythematosus


MW: 12 kDa

**Gene Summary:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. Several transcript variants encoding different isoforms, have been identified for this gene. [provided by RefSeq, Oct 2015]

### Product images:



Circular map for RC217665



Western blot validation of overexpression lysate (Cat# [LY408460]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC217665 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).