

Product datasheet for RC217669

MRP5 (ABCC5) (NM_001023587) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: MRP5 (ABCC5) (NM_001023587) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: MRP5
Synonyms: ABC33; EST277145; MOAT-C; MOATC; MRP5; pABC11; SMRP
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC217669 representing NM_001023587
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAAGGATATCGACATAGGAAAAGAGTATATCATCCCCAGTCTGGGTATAGAAGTGTGAGGGAGAGAA
CCAGCACTTCTGGGACGCACAGAGACCGTGAAGATTCCAAGTTCAGGAGAACTCGACCGTTGGAATGCCA
AGATGCCTTGAAACAGCAGCCCGAGCCGAGGGCCTCTCTTTGATGCCTCCATGCATTCTCAGCTCAGA
ATCCTGGATGAGGAGCATCCAAGGAAAGTACCATCATGGCTTGAGTGTCTGAAGCCCATCCGGACTA
CTTCCAAACACCAGCACCCAGTGGACAATGCTGGGCTTTTTTCCGTATGACTTTTTCTGGCTTTCTTC
TCTGGCCCGTGTGGCCCAAGAAGGGGAGCTCTCAATGGAAGACGTGTGGTCTCTGTCCAAGCAGAG
TCTTCTGACGTGAAGTGCAGAAGACTAGAGAGACTGTGGCAAGAAGAGCTGAATGAAGTTGGCCAGACG
CTGCTTCCCTGCGAAGGGTTGTGTGGATCTTCTGCCGACCAGGCTCATCCTGTCCATCGTGTGCCTGAT
GATCACGCAGCTGGCTGGCTTCACTGGACCAAATTTTCAGGATGGCTGTATTCTGCGGTCAGAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC217669 representing NM_001023587
Red=Cloning site Green=Tags(s)

MKDIDIGKEYIIPSPGYRSVRERTSTSGTHRDREDSKFRTRRPLECQDALETAARAEGLSLDASMHSQLR
ILDEEHPKGKYHHGLSALKPIRTTSKHQHPVDNAGLFSMTFSWLSSLARVAHKKGELSMEDVWLSKHE
SSDVNCRRLERLWQEELNEVGPDAASLRVWVIFCRTRLILSIVCLMITQLAGFSGPNFQDGCILRSE


TRTRPLEQKLISEEDLAANDILDYKDDDDKV



Chromatograms: https://cdn.origene.com/chromatograms/mk8004_g02.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001023587

ORF Size: 624 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001023587.3
RefSeq Size:	2007 bp
RefSeq ORF:	627 bp
Locus ID:	10057
UniProt ID:	O15440
Cytogenetics:	3q27.1
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	ABC transporters
MW:	23.5 kDa
Gene Summary:	<p>The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions in the cellular export of its substrate, cyclic nucleotides. This export contributes to the degradation of phosphodiesterases and possibly an elimination pathway for cyclic nucleotides. Studies show that this protein provides resistance to thiopurine anticancer drugs, 6-mercaptopurine and thioguanine, and the anti-HIV drug 9-(2-phosphonylmethoxyethyl)adenine. This protein may be involved in resistance to thiopurines in acute lymphoblastic leukemia and antiretroviral nucleoside analogs in HIV-infected patients. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]</p>


Product images:



Circular map for RC217669



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ABCC5 (Cat# RC217669, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ABCC5 (Cat# [TA808487])(1:2000). Positive lysates [LY422609] (100ug) and [LC422609] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY422609]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC217669 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).