

Product datasheet for RC217737L4

QDPR (NM_000320) Human Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	QDPR (NM_000320) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	QDPR
Synonyms:	DHPR; HDHPR; PKU2; SDR33C1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC217737).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I ORF Mlu I [GCG ATC GC] ATG // NNN ACG CGT]



ACCN: ORF Size: NM_000320 732 bp



View online »


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

QDPR (NM_000320) Human Tagged Lenti ORF Clone – RC217737L4

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailin variants is recommended prior to use. More infoOTI Annotation:This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of waterReconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liqu at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.RefSeq:NM 000320.1RefSeq ORF:735 bpLocus ID:909417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene	OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of waterReconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liqu at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date or shipping when stored at -20°C.RefSeq:NM 000320.1RefSeq ORF:735 bpLocus ID:5860UniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine		reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing
containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of waterReconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liqu at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.RefSeq:NM 000320.1RefSeq ORF:735 bpLocus ID:5860UniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- 	OTI Annotation:	
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liqu at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date or shipping when stored at -20°C. RefSeq: NM 000320.1 RefSeq ORF: 735 bp Locus ID: 5860 UniProt ID: P09417 Cytogenetics: 4p15.32 Protein Families: Druggable Genome Protein Pathways: Folate biosynthesis, Metabolic pathways MW: 25.6 kDa Gene Summary: This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH-mediated reduction of quinonoid dihydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine 	Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq Size:1550 bpRefSeq ORF:735 bpLocus ID:5860UniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	Reconstitution Method:	 Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of
RefSeq ORF:735 bpLocus ID:5860UniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	RefSeq:	<u>NM 000320.1</u>
Locus ID:5860UniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysWW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH-mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	RefSeq Size:	1550 bp
JniProt ID:P09417Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysVW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	RefSeq ORF:	735 bp
Cytogenetics:4p15.32Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	ocus ID:	5860
Protein Families:Druggable GenomeProtein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	JniProt ID:	<u>P09417</u>
Protein Pathways:Folate biosynthesis, Metabolic pathwaysMW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	Cytogenetics:	4p15.32
MW:25.6 kDaGene Summary:This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	Protein Families:	Druggable Genome
Gene Summary: This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH- mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	Protein Pathways:	Folate biosynthesis, Metabolic pathways
mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine	/W:	25.6 kDa
	Gene Summary:	mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component of the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, or premature terminations. Dihydropteridine reductase deficiency presents as atypical phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalanine

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Circular map for RC217737L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US