

## Product datasheet for RC217864L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RTN3 (NM 201429) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** RTN3 (NM\_201429) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ASYIP; HAP; NSPL2; NSPLII; RTN3-A1 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 201429

**ORF Size:** 765 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC217864).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

NM 201429.2, NP 958832.1 RefSeq:

RefSeq Size: 2645 bp RefSeq ORF: 768 bp Locus ID: 10313 **UniProt ID:** O95197 Cytogenetics: 11q13.1

**Protein Families:** Transmembrane

MW: 27.6 kDa







## **Gene Summary:**

This gene belongs to the reticulon family of highly conserved genes that are preferentially expressed in neuroendocrine tissues. This family of proteins interact with, and modulate the activity of beta-amyloid converting enzyme 1 (BACE1), and the production of amyloid-beta. An increase in the expression of any reticulon protein substantially reduces the production of amyloid-beta, suggesting that reticulon proteins are negative modulators of BACE1 in cells. Alternatively spliced transcript variants encoding different isoforms have been found for this gene, and pseudogenes of this gene are located on chromosomes 4 and 12. [provided by RefSeq, May 2012]