

## Product datasheet for RC217876L3

### LARGE (LARGE1) (NM\_004737) Human Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	LARGE (LARGE1) (NM_004737) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	LARGE
Synonyms:	LARGE; MDC1D; MDDGA6; MDDGB6
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC217876).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_004737
ORF Size:	2268 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_004737.3</a>
<b>RefSeq Size:</b>	4194 bp
<b>RefSeq ORF:</b>	2271 bp
<b>Locus ID:</b>	9215
<b>UniProt ID:</b>	<a href="#">O95461</a>
<b>Cytogenetics:</b>	22q12.3
<b>Domains:</b>	Glyco_transf_8
<b>Protein Families:</b>	Druggable Genome, Transmembrane
<b>MW:</b>	87.9 kDa
<b>Gene Summary:</b>	This gene encodes a member of the N-acetylglucosaminyltransferase gene family. It encodes a glycosyltransferase which participates in glycosylation of alpha-dystroglycan, and may carry out the synthesis of glycoprotein and glycosphingolipid sugar chains. It may also be involved in the addition of a repeated disaccharide unit. The protein encoded by this gene is the glycotransferase that adds the final xylose and glucuronic acid to alpha-dystroglycan and thereby allows alpha-dystroglycan to bind ligands including laminin 211 and neurexin. Mutations in this gene cause several forms of congenital muscular dystrophy characterized by cognitive disability and abnormal glycosylation of alpha-dystroglycan. Alternative splicing of this gene results in multiple transcript variants that encode the same protein. [provided by RefSeq, May 2018]

Product images:



Circular map for RC217876L3