

Product datasheet for RC217954L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Repulsive Guidance Molecule C (HFE2) (NM_213653) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Repulsive Guidance Molecule C (HFE2) (NM_213653) Human Tagged ORF Clone Lentiviral

Particle

Symbol: HJV

Synonyms: HFE2; HFE2A; JH; RGMC

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_213653 **ORF Size:** 1278 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC217954).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 213653.2</u>

 RefSeq Size:
 2128 bp

 RefSeq ORF:
 1281 bp

 Locus ID:
 148738

 UniProt ID:
 Q6ZVN8

 Cytogenetics:
 1q21.1

Protein Families: Transmembrane





MW: 41.5 kDa

Gene Summary: The product of this gene is involved in iron metabolism. It may be a component of the

signaling pathway which activates hepcidin or it may act as a modulator of hepcidin expression. It could also represent the cellular receptor for hepcidin. Two uORFs in the 5' UTR negatively regulate the expression and activity of the encoded protein. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. Defects in this gene are the cause of hemochromatosis type 2A, also called juvenile hemochromatosis (JH). JH is an early-onset autosomal recessive disorder due to severe iron overload resulting in hypogonadotrophic hypogonadism, hepatic fibrosis or cirrhosis and cardiomyopathy,

occurring typically before age of 30. [provided by RefSeq, Oct 2015]