

Product datasheet for RC218040L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SHIP (INPP5D) (NM_001017915) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: SHIP (INPP5D) (NM_001017915) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: SHIP

Synonyms: hp51CN; p150Ship; SHIP; SHIP-1; SHIP1; SIP-145

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC218040).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001017915

ORF Size: 3567 bp



SHIP (INPP5D) (NM_001017915) Human Tagged Lenti ORF Clone - RC218040L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001017915.1</u>, <u>NP 001017915.1</u>

2q37.1

 RefSeq Size:
 4928 bp

 RefSeq ORF:
 3570 bp

 Locus ID:
 3635

 UniProt ID:
 Q92835

Cytogenetics:

Protein Families: Druggable Genome

Protein Pathways: B cell receptor signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated

phagocytosis, Insulin signaling pathway, Phosphatidylinositol signaling system

MW: 133.1 kDa

Gene Summary: This gene is a member of the inositol polyphosphate-5-phosphatase (INPP5) family and

encodes a protein with an N-terminal SH2 domain, an inositol phosphatase domain, and two

C-terminal protein interaction domains. Expression of this protein is restricted to hematopoietic cells where its movement from the cytosol to the plasma membrane is mediated by tyrosine phosphorylation. At the plasma membrane, the protein hydrolyzes the

5' phosphate from phosphatidylinositol (3,4,5)-trisphosphate and inositol-1,3,4,5-

tetrakisphosphate, thereby affecting multiple signaling pathways. The protein is also partly localized to the nucleus, where it may be involved in nuclear inositol phosphate signaling processes. Overall, the protein functions as a negative regulator of myeloid cell proliferation and survival. Mutations in this gene are associated with defects and cancers of the immune system. Deficiencies in the encoded protein, SHIP1, have been associated with Inflammatory Bowel Disease types such as Crohn's Disease and Ulcerative Colitis. Alternative splicing of this

gene results in multiple transcript variants. [provided by RefSeq, Jul 2020]



Product images:

