

Product datasheet for RC218134L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SIRT1 (NM_012238) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SIRT1 (NM_012238) Human Tagged ORF Clone Lentiviral Particle

Symbol: SIRT'

Synonyms: SIR2; SIR2alpha; SIR2L1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 012238

ORF Size: 2241 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC218134).

Sequence:

, , ,

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

The molecular sequence of this clone aligns with the gene accession number as a point of

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 012238.3

 RefSeq Size:
 4107 bp

 RefSeq ORF:
 2244 bp

 Locus ID:
 23411

 UniProt ID:
 Q96EB6

 Cytogenetics:
 10q21.3

Domains: SIR2

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors





ORIGENE

MW: 81.5 kDa

Gene Summary:

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2008]