

## Product datasheet for RC218299L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NFAT5 (NM\_173214) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** NFAT5 (NM\_173214) Human Tagged ORF Clone Lentiviral Particle

Symbol: NFAT5

Synonyms: NF-AT5; NFATL1; NFATZ; OREBP; TONEBP

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_173214

 ORF Size:
 4647 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC218299).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 173214.1</u>

 RefSeq Size:
 13315 bp

 RefSeq ORF:
 4368 bp

 Locus ID:
 10725

 UniProt ID:
 094916

Cytogenetics: 16q22.1

**Protein Families:** Druggable Genome, Transcription Factors





## NFAT5 (NM\_173214) Human Tagged ORF Clone Lentiviral Particle - RC218299L3V

**Protein Pathways:** Axon guidance, B cell receptor signaling pathway, Natural killer cell mediated cytotoxicity, T

cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway

**MW:** 167.7 kDa

**Gene Summary:** The product of this gene is a member of the nuclear factors of activated T cells family of

transcription factors. Proteins belonging to this family play a central role in inducible gene transcription during the immune response. This protein regulates gene expression induced by osmotic stress in mammalian cells. Unlike monomeric members of this protein family, this protein exists as a homodimer and forms stable dimers with DNA elements. Multiple

transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008]