

# Product datasheet for RC218310L3V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PSMA (FOLH1) (NM\_004476) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** PSMA (FOLH1) (NM\_004476) Human Tagged ORF Clone Lentiviral Particle

Symbol: PSMA

Synonyms: FGCP; FOLH; GCP2; GCPII; mGCP; NAALAD1; NAALAdase; PSM; PSMA

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM\_004476

ORF Size: 2250 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC218310).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 004476.1

 RefSeq Size:
 2653 bp

 RefSeq ORF:
 2253 bp

 Locus ID:
 2346

 UniProt ID:
 Q04609

 Cytogenetics:
 11p11.12

**Domains:** PA, TFR\_dimer, Peptidase\_M28

**Protein Families:** Druggable Genome, Protease, Transmembrane






MW:

84.2 kDa

#### **Gene Summary:**


This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase family. The protein acts as a glutamate carboxypeptidase on different alternative substrates, including the nutrient folate and the neuropeptide N-acetyl-l-aspartyl-l-glutamate and is expressed in a number of tissues such as prostate, central and peripheral nervous system and kidney. A mutation in this gene may be associated with impaired intestinal absorption of dietary folates, resulting in low blood folate levels and consequent hyperhomocysteinemia. Expression of this protein in the brain may be involved in a number of pathological conditions associated with glutamate excitotoxicity. In the prostate the protein is upregulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal region. Alternative splicing gives rise to multiple transcript variants encoding several different isoforms. [provided by RefSeq, Jul 2010]

## **Product images:**



[RC218310L3] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC218310L3V particle to overexpress human FOLH1-Myc-DDK fusion protein.





[RC218310L3] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC218310L3V particle to overexpress human FOLH1-Myc-DDK fusion protein.