

Product datasheet for **RC218439**

cGKI (PRKG1) (NM_001098512) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	cGKI (PRKG1) (NM_001098512) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	cGKI
Synonyms:	AAT8; cGK; cGK 1; cGK1; cGKI; cGKI-alpha; cGKI-BETA; PKG; PKG1; PRKG1B; PRKGR1B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RC218439 representing NM_001098512
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCGCGATCGCC

ATGAGCGAGCTAGAGGAAGACTTTGCCAAGATTCTCATGCTCAAGGAGGAGAGGATCAAAGAGCTGGAGA
 AGCGGCTGTCAGAGAAGGAGGAAGAAATTCAGGAGCTGAAGAGGAACTCCACAAATGCCAGTCGGTGCT
 CCCAGTGCCCTCGACCCACATCGGCCCCCGGACCACCCGGGCGCAGGGCATCTCGGCCGAGCCGACAGC
 TACAGGTCCTTCCACGACCTCCGACAGGCATTCCGGAAGTTCACCAAGTCCGAAAGGTCCAAGGATCTTA
 TAAAGGAAGCTATCCTTGACAATGACTTTATGAAGAACTTGGAGCTGTCGAGATCCAGGAGATTGTGGA
 TTGTATGTACCCGGTGGAGTATGGCAAGGACAGTTGCATCATCAAAGAAGGAGACGTGGGGTCACTGGTG
 TATGTCATGGAAGATGGTAAGGTTGAAGTTACAAAAGAAGGTGTGAAGTTGTGTACCATGGTCCAGGAA
 AAGTGTGGGGAATTGGCTATTCTTTACAACGTACCCGGACAGCGACCGTCAAGACTCTGTAATGT
 AAAACTCTGGGCCATTGATCGACAATGTTTTCAAACAATAATGATGAGGACAGGACTCATCAAGCATAACC
 GAGTATATGGAATTTTTAAAAAGCGTTCCAACATTCCAGAGCCTTCTGAAGAGATCCTCAGCAAGCTTG
 CTGATGTCCTTGAAGAGACCCACTATGAAAATGGAGAATATATTATCAGGCAAGGTGCAAGAGGGGACAC
 CTTCTTTATCATCAGCAAAGGAACGGTAAATGTCACTCGTGAAGACTCACCGAGTGAAGACCCAGTCTTT
 CTTAGAACTTTAGGAAAAGGAGACTGGTTTGGAGAGAAAGCCTTGCAGGGGGAAGATGTGAGAACAGCAA
 ACGTAATTGCTGCAGAAGCTGTAACTGCCTTGTGATTGACAGAGACTCTTTAAACATTTGATTGGAGG
 GCTGGATGATGTTCTAATAAAGCATATGAAGATGCAGAAGCTAAAGCAAAATATGAAGCTGAAGCGGCT
 TTCTTCGCCAACCTGAAGCTGTCTGATTCAACATCATTGATACCCTTGGAGTTGGAGGTTTCGGACGAG
 TAGAACTGGTCCAGTTGAAAAGTGAAGAATCCAAAACGTTTGAATGAAGATTCTCAAGAAACGTCACAT
 TGTGGACACAAGACAGCAGGAGCACATCCGCTCAGAGAAGCAGATCATGCAGGGGGCTCATTCCGATTTT
 ATAGTGAGACTGTACAGAACATTTAAGGACAGCAAATATTTGTATATGTTGATGGAAGCTTGTCTAGGTG
 GAGAGCTCTGGACCATTCTCAGGGATAGAGGTTTCGTTTGAAGATTCTACAACCAGATTTTACACAGCATG
 TGTGGTAGAAGCTTTTGCCTATCTGCATTCCAAAGGAATCATTTACAGGGACCTCAAGCCAGAAAATCTC
 ATCCTAGATCACCGAGGTTATGCCAACTGGTTGATTTTGGCTTTGCAAAGAAAATAGGATTTGAAAGA
 AAACATGGACTTTTTGTGGGACTCCAGAGTATGTAGCCCCAGAGATCATCCTGAACAAAGGCCATGACAT
 TTCAGCCGACTACTGGTCACTGGGAATCCTAATGTATGAACTCCTGACTGGCAGCCACCTTTCAGGC
 CCAGATCCTATGAAAACCTATAACATCATATTGAGGGGATTGACATGATAGAATTTCCAAGAAGATTG
 CCAAAAATGCTGCTAATTTAATTAATAAAAACTATGCAGGGACAATCCATCAGAAAAGATTAGGGAATTTGAA
 AAATGGAGTAAAAGACATTCAAAGCACAATGGTTTGAAGGCTTAACTGGGAAGGCTTAAAGAAAAGGT
 ACCTTGACACCTCCTATAATACCAAGTGTTCATCACCCACAGACACAAGTAATTTTACAGTTCCTG
 AGGACAACGATGAACCACCACCTGATGACAACCTCAGGATGGGATATAGACTTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001098512.3](#)

RefSeq Size: 3824 bp

RefSeq ORF: 2016 bp

Locus ID: 5592

UniProt ID: [Q13976](#)

Cytogenetics: 10q11.23-q21.1


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Gap junction, Long-term depression, Olfactory transduction, Vascular smooth muscle contraction


MW: 76.2 kDa

Gene Summary: Mammals have three different isoforms of cyclic GMP-dependent protein kinase (Ialpha, Ibeta, and II). These PRKG isoforms act as key mediators of the nitric oxide/cGMP signaling pathway and are important components of many signal transduction processes in diverse cell types. This PRKG1 gene on human chromosome 10 encodes the soluble Ialpha and Ibeta isoforms of PRKG by alternative transcript splicing. A separate gene on human chromosome 4, PRKG2, encodes the membrane-bound PRKG isoform II. The PRKG1 proteins play a central role in regulating cardiovascular and neuronal functions in addition to relaxing smooth muscle tone, preventing platelet aggregation, and modulating cell growth. This gene is most strongly expressed in all types of smooth muscle, platelets, cerebellar Purkinje cells, hippocampal neurons, and the lateral amygdala. Isoforms Ialpha and Ibeta have identical cGMP-binding and catalytic domains but differ in their leucine/isoleucine zipper and autoinhibitory sequences and therefore differ in their dimerization substrates and kinase enzyme activity. [provided by RefSeq, Sep 2011]

Product images:



Circular map for RC218439



Western blot validation of overexpression lysate (Cat# [LY420611]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC218439 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).