

## Product datasheet for RC218519L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Estrogen Receptor beta (ESR2) (NM\_001437) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Estrogen Receptor beta (ESR2) (NM\_001437) Human Tagged ORF Clone Lentiviral Particle

Symbol: Estrogen Receptor beta

**Synonyms:** ER-BETA; Erb; ESR-BETA; ESRB; ESTRB; NR3A2; ODG8

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 001437

ORF Size: 1590 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC218519).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001437.1

RefSeq Size: 2011 bp
RefSeq ORF: 1593 bp
Locus ID: 2100

 UniProt ID:
 Q92731

 Cytogenetics:
 14q23.2-q23.3

Cytogenetics: 14q23.2-q23

Domains: HOLI, zf-C4

**Protein Families:** Druggable Genome, Nuclear Hormone Receptor, Transcription Factors





Estrogen Receptor beta (ESR2) (NM\_001437) Human Tagged ORF Clone Lentiviral Particle – RC218519L3V

MW:

59 kDa

**Gene Summary:** 

This gene encodes a member of the family of estrogen receptors and superfamily of nuclear receptor transcription factors. The gene product contains an N-terminal DNA binding domain and C-terminal ligand binding domain and is localized to the nucleus, cytoplasm, and mitochondria. Upon binding to 17beta-estradiol or related ligands, the encoded protein forms homo- or hetero-dimers that interact with specific DNA sequences to activate transcription. Some isoforms dominantly inhibit the activity of other estrogen receptor family members. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been fully characterized. [provided by RefSeq, Jul 2008]