

Product datasheet for RC218694

RARRES1 (NM_002888) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	RARRES1 (NM_002888) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	RARRES1
Synonyms:	LXNL; PERG-1; TIG1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC218694 representing NM_002888 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAAGGGCGCCGGGAATTCTCGACTGGATCCGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGCAGCCCCGCCGCAACGGCTGCCTGCCCTGGTCCGGCCCAGGGGCCGCCACCGCCCCGC
 TGCTCGCCTGCTGCTGTTGCTGCCCGTGGCGGCCGCCGGGTCCGGGGACCCGACGACCCCTGG
 GCAGCCTCAGGATGCTGGGTCCCGCGCAGGCTCCTGCAGCAGGCCGCCGCCGCCGCTTCACTTCTC
 AACTTCCGGTCCGGCTGCCAGCGCGCTACGAGTGCTGGCCGAGGTGCAGGAGGCCGCCGTGGATT
 ATCCAAAAGAGGGATGTAAGTTCACGTGGTCTTCAGCACAGAGCGCTACAACCCAGAGTCTTACTTCA
 GGAAGGTAGGGACGTTGGGAAATGTTCTGCTCGAGTGTTTCAAGAATCAGAAACCCAGACCAACC
 ATCAATGTAACTGTACACGGCTCATCGAGAAAAAGAAAAGACAACAAGAGGATTACCTGCTTTACAAGC
 AAATGAAGCACTGAAAACCCCTTGGAAATAGTCAGCATACCTGATAATCATGGACATATTGATCCCTC
 TCTGAGACTCATGGATTGGCTTCTGGAAAGCTTACGTGATGTGGAAATGACAACACAGTG
 TCACACTACTTGGCACAGCTACTAGTGTGAGGCAGTGGTAAGAAAACC

ACCGTACGCGCCGCTCGAGCAGAAACTCATCTCAGAACAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTAA



[View online »](#)



Protein Sequence: >RC218694 representing NM_002888
 Red=Cloning site Green=Tags(s)

MQPQQRQLPAPWSGPRGPRPTAPLLALLLAPVAAPAGSGDPDPGPQDAGVPRRLQQAARAALHFF
 NFRSGSPSALRVLAEVQEGRRAWINPKEGCKVHVFSTERYNPESLLQEGERGLGKCSARVFFKNQKPRPT
 INVTCCTRLIEKKRQQEDYLLYKQMKQLKNPLEIVSIPDNHGHDPSLRIWDLAFLGSSYVMEMTTQV
 SHYLAQLTSVRQWRKT

TRTRPLEEQKLISEEDLAANDILDYKDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:



* The last codon before the Stop codon of the ORF

ACCN: NM_002888


ORF Size: 684 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_002888.3
RefSeq Size:	886 bp
RefSeq ORF:	687 bp
Locus ID:	5918
UniProt ID:	P49788
Cytogenetics:	3q25.32
Protein Families:	Druggable Genome
MW:	25.6 kDa
Gene Summary:	This gene was identified as a retinoid acid (RA) receptor-responsive gene. It encodes a type 1 membrane protein. The expression of this gene is upregulated by tazarotene as well as by retinoic acid receptors. The expression of this gene is found to be downregulated in prostate cancer, which is caused by the methylation of its promoter and CpG island. Alternatively spliced transcript variant encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]

Product images:

Circular map for RC218694