

Product datasheet for RC218905L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

beta TRCP2 (FBXW11) (NM_012300) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: beta TRCP2 (FBXW11) (NM_012300) Human Tagged ORF Clone Lentiviral Particle

Symbol: beta TRCP2

Synonyms: BTRC2; BTRCP2; FBW1B; Fbw11; FBXW1B; Hos; NEDJED

Mammalian Cell

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

NM 012300

Tag: Myc-DDK

ORF Size: 1626 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC218905).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 012300.2</u>

RefSeq Size: 4575 bp
RefSeq ORF: 1629 bp
Locus ID: 23291
UniProt ID: Q9UKB1
Cytogenetics: 5q35.1

Domains: WD40, F-box

Protein Families: Druggable Genome





beta TRCP2 (FBXW11) (NM_012300) Human Tagged ORF Clone Lentiviral Particle - RC218905L1V

Protein Pathways: Hedgehog signaling pathway, Oocyte meiosis, Ubiquitin mediated proteolysis, Wnt signaling

pathway

MW: 61.9 kDa

Gene Summary: This gene encodes a member of the F-box protein family which is characterized by an

approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbws class and, in addition to an F-box, contains multiple WD40 repeats. This gene contains at least 14 exons, and its alternative splicing generates 3 transcript variants diverging at the presence/absence of two alternate exons. [provided by

RefSeq, Jul 2008]