

Product datasheet for RC218955L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CEBP Alpha (CEBPA) (NM 004364) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CEBP Alpha (CEBPA) (NM 004364) Human Tagged ORF Clone Lentiviral Particle

Symbol: CEBP Alpha

Synonyms: C/EBP-alpha; CEBP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 004364

ORF Size: 1074 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC218955).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally accurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004364.2

 RefSeq Size:
 2385 bp

 RefSeq ORF:
 1077 bp

 Locus ID:
 1050

 UniProt ID:
 P49715

Cytogenetics: 19q13.11

Protein Families: Druggable Genome, ES Cell Differentiation/IPS

Protein Pathways: Acute myeloid leukemia, Pathways in cancer






MW:

37.4 kDa

Gene Summary:

This intronless gene encodes a transcription factor that contains a basic leucine zipper (bZIP) domain and recognizes the CCAAT motif in the promoters of target genes. The encoded protein functions in homodimers and also heterodimers with CCAAT/enhancer-binding proteins beta and gamma. Activity of this protein can modulate the expression of genes involved in cell cycle regulation as well as in body weight homeostasis. Mutation of this gene is associated with acute myeloid leukemia. The use of alternative in-frame non-AUG (GUG) and AUG start codons results in protein isoforms with different lengths. Differential translation initiation is mediated by an out-of-frame, upstream open reading frame which is located between the GUG and the first AUG start codons. [provided by RefSeq, Dec 2013]

Product images:



[RC218955L3] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC218955L3V particle to overexpress human CEBPA-Myc-DDK fusion protein.