

Product datasheet for RC218992L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SRC3 (NCOA3) (NM_006534) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SRC3 (NCOA3) (NM_006534) Human Tagged ORF Clone Lentiviral Particle

Symbol: SRC3

Synonyms: ACTR; AIB-1; AIB1; bHLHe42; CAGH16; CTG26; KAT13B; pCIP; RAC3; SRC-3; SRC3; TNRC14;

TNRC16; TRAM-1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_006534

ORF Size: 4260 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC218992).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006534.2</u>

 RefSeq Size:
 7923 bp

 RefSeq ORF:
 4263 bp

 Locus ID:
 8202

 UniProt ID:
 Q9Y6Q9

 Cytogenetics:
 20q13.12

Domains: PAS, HLH





SRC3 (NCOA3) (NM_006534) Human Tagged ORF Clone Lentiviral Particle - RC218992L3V

Protein Families: Druggable Genome, Transcription Factors

MW: 154.7 kDa

Gene Summary: The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear

hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript

variants encoding different isoforms have been found for this gene. In addition, a

polymorphic repeat region is found in the C-terminus of the encoded protein. [provided by

RefSeq, Mar 2010]