

## Product datasheet for RC219000L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TAC4 (NM\_001077505) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** TAC4 (NM\_001077505) Human Tagged ORF Clone Lentiviral Particle

Symbol: TAC4

Synonyms: EK; HK-1; HK1; PPT-C

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001077505

ORF Size: 228 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219000).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001077505.1, NP 001070973.1</u>

 RefSeq Size:
 564 bp

 RefSeq ORF:
 231 bp

 Locus ID:
 255061

 UniProt ID:
 Q86UU9

 Cytogenetics:
 17q21.33

 MW:
 7.8 kDa







## **Gene Summary:**

This gene is a member of the tachykinin family of neurotransmitter-encoding genes. Tachykinin proteins are cleaved into small, secreted peptides that activate members of a family of receptor proteins. The products of this gene preferentially activate tachykinin receptor 1, and are thought to regulate peripheral endocrine and paracrine functions including blood pressure, the immune system, and endocrine gland secretion. The products of this gene lack a dibasic cleavage site found in other tachykinin proteins. Consequently, the nature of the cleavage products generated in vivo remains to be determined. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]