

## Product datasheet for RC219027L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL29 (IFNL1) (NM\_172140) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** IL29 (IFNL1) (NM\_172140) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL29

Synonyms: IL-29; IL29

Mammalian Cell None

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 172140

ORF Size: 600 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219027).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 172140.1

 RefSeq Size:
 856 bp

 RefSeq ORF:
 603 bp

 Locus ID:
 282618

 UniProt ID:
 Q8IU54

 Cytogenetics:
 19q13.2

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway





ORIGENE

**MW:** 21.9 kDa

**Gene Summary:** 

This gene encodes a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 28B (IL28B) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA). [provided by RefSeq, Jul 2008]