

## Product datasheet for RC219033L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TFPI (NM 006287) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** TFPI (NM\_006287) Human Tagged ORF Clone Lentiviral Particle

Symbol:

EPI; LACI; TFI; TFPI1 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

NM 006287 ACCN:

**ORF Size:** 912 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC219033).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 006287.3

RefSeq Size: 1431 bp RefSeq ORF: 915 bp Locus ID: 7035 **UniProt ID:** P10646 Cytogenetics: 2q32.1

**Domains:** KIJ

**Protein Families:** Secreted Protein





## TFPI (NM\_006287) Human Tagged ORF Clone Lentiviral Particle - RC219033L4V

**Protein Pathways:** Complement and coagulation cascades

MW: 35.02 kDa

Gene Summary: This gene encodes a Kunitz-type serine protease inhibitor that regulates the tissue factor (TF)-

dependent pathway of blood coagulation. The coagulation process initiates with the formation of a factor VIIa-TF complex, which proteolytically activates additional proteases (factors IX and X) and ultimately leads to the formation of a fibrin clot. The product of this gene inhibits the activated factor X and VIIa-TF proteases in an autoregulatory loop. Inhibition of the encoded protein restores hemostasis in animal models of hemophilia. This gene encodes multiple protein isoforms that differ in their inhibitory activity, specificity and cellular

localization. [provided by RefSeq, Jul 2016]