

Product datasheet for RC219169L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CHIT1 (NM_003465) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CHIT1 (NM_003465) Human Tagged ORF Clone Lentiviral Particle

Symbol: CHIT1

Synonyms: CHI3; CHIT; CHITD

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag:Myc-DDKACCN:NM_003465

ORF Size: 1398 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219169).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003465.1

RefSeq Size: 1633 bp
RefSeq ORF: 1401 bp
Locus ID: 1118
UniProt ID: Q13231
Cytogenetics: 1q32.1

Domains: Glyco_18, CBM_14

Protein Families: Secreted Protein, Transmembrane





CHIT1 (NM_003465) Human Tagged ORF Clone Lentiviral Particle - RC219169L1V

Protein Pathways: Amino sugar and nucleotide sugar metabolism

MW: 51.5 kDa

Gene Summary: Chitotriosidase is secreted by activated human macrophages and is markedly elevated in

plasma of Gaucher disease patients. The expression of chitotriosidase occurs only at a late

stage of differentiation of monocytes to activated macrophages in culture. Human

macrophages can synthesize a functional chitotriosidase, a highly conserved enzyme with a strongly regulated expression. This enzyme may play a role in the degradation of chitin-containing pathogens. Several alternatively spliced transcript variants have been described

for this gene. [provided by RefSeq, Jan 2012]