

Product datasheet for RC219172L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDGFC (NM_016205) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PDGFC (NM_016205) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDGFC

Synonyms: FALLOTEIN; SCDGF

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 016205

Tag: Myc-DDK

ORF Size: 1035 bp

ORF Nucleotide

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC219172).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 016205.1

 RefSeq Size:
 3007 bp

 RefSeq ORF:
 1038 bp

 Locus ID:
 56034

 UniProt ID:
 Q9NRA1

Cytogenetics: 4q32.1

Domains: PDGF, CUB



PDGFC (NM_016205) Human Tagged ORF Clone Lentiviral Particle - RC219172L3V

Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,

Induced pluripotent stem cells

Protein Pathways: Cytokine-cytokine receptor interaction, Focal adhesion, Gap junction, Melanoma, Prostate

cancer, Regulation of actin cytoskeleton

MW: 39.03 kDa

Gene Summary: The protein encoded by this gene is a member of the platelet-derived growth factor family.

The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a core motif of eight cysteines. This gene product appears to form only homodimers. It differs from the platelet-derived growth factor alpha and beta polypeptides in

having an unusual N-terminal domain, the CUB domain. Alternatively spliced transcript

variants have been found for this gene. [provided by RefSeq, Sep 2010]