

## Product datasheet for RC219253L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **HCST (NM\_001007469) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** HCST (NM\_001007469) Human Tagged ORF Clone Lentiviral Particle

Symbol: HCST

Synonyms: DAP10; KAP10; PIK3AP

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001007469

ORF Size: 279 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219253).

Sequence:

Cytogenetics:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001007469.1, NP 001007470.1

RefSeq Size:521 bpRefSeq ORF:279 bpLocus ID:10870UniProt ID:Q9UBK5

**Protein Families:** Druggable Genome, Transmembrane

19q13.12

**Protein Pathways:** Natural killer cell mediated cytotoxicity





ORIGENE

**MW:** 9.5 kDa

**Gene Summary:** 

This gene encodes a transmembrane signaling adaptor that contains a YxxM motif in its cytoplasmic domain. The encoded protein may form part of the immune recognition receptor complex with the C-type lectin-like receptor NKG2D. As part of this receptor complex, this protein may activate phosphatidylinositol 3-kinase dependent signaling pathways through its intracytoplasmic YxxM motif. This receptor complex may have a role in cell survival and proliferation by activation of NK and T cell responses. Alternative splicing results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]