

Product datasheet for RC219265L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SIGLECL1 (SIGLEC12) (NM_033329) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: SIGLECL1 (SIGLEC12) (NM_033329) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: SIGLECL1

Synonyms: S2V; Siglec-XII; SIGLECL1; SLG

Puromycin

Mammalian Cell

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of the

Sequence:

The ORF insert of this clone is exactly the same as(RC219265).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_033329

ORF Size: 1431 bp





SIGLECL1 (SIGLEC12) (NM_033329) Human Tagged Lenti ORF Clone - RC219265L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 033329.1, NP 201586.1

 RefSeq Size:
 1736 bp

 RefSeq ORF:
 1434 bp

 Locus ID:
 89858

 UniProt ID:
 Q96PQ1

 Cytogenetics:
 19q13.41

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transmembrane

MW: 50.1 kDa

Gene Summary: Sialic acid-binding immunoglobulin-like lectins (SIGLECs) are a family of cell surface proteins

belonging to the immunoglobulin superfamily. They mediate protein-carbohydrate


interactions by selectively binding to different sialic acid moieties present on glycolipids and glycoproteins. This gene encodes a member of the SIGLEC3-like subfamily of SIGLECs. Members of this subfamily are characterized by an extracellular V-set immunoglobulin-like domain followed by two C2-set immunoglobulin-like domains, and the cytoplasmic tyrosine-

based motifs ITIM and SLAM-like. The encoded protein, upon tyrosine phosphorylation, has been shown to recruit the Src homology 2 domain-containing protein-tyrosine phosphatases SHP1 and SHP2. It has been suggested that the protein is involved in the negative regulation of macrophage signaling by functioning as an inhibitory receptor. This gene is located in a cluster with other SIGLEC3-like genes on 19q13.4. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Aug 2013]



Product images:



Circular map for RC219265L4