

Product datasheet for RC219275L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Delta Opioid Receptor (OPRD1) (NM_000911) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Delta Opioid Receptor (OPRD1) (NM_000911) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Delta Opioid Receptor

Synonyms: DOP; DOR; DOR1; OPRD

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC219275).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_000911

ORF Size: 1116 bp





Delta Opioid Receptor (OPRD1) (NM_000911) Human Tagged Lenti ORF Clone - RC219275L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000911.3</u>

RefSeq Size:1774 bpRefSeq ORF:1119 bpLocus ID:4985

UniProt ID: P41143

Cytogenetics: 1p35.3

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Neuroactive ligand-receptor interaction

MW: 40.2 kDa

Gene Summary: G-protein coupled receptor that functions as receptor for endogenous enkephalins and for a


subset of other opioids. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase. Signaling leads to the inhibition of adenylate cyclase activity. Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Plays a role in the perception of pain and in opiate-

mediated analgesia. Plays a role in developing analgesic tolerance to morphine.


[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for RC219275L1



Double digestion of RC219275L1 using Sgfl and Mlul