

Product datasheet for RC219275L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Delta Opioid Receptor (OPRD1) (NM 000911) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Delta Opioid Receptor (OPRD1) (NM_000911) Human Tagged ORF Clone Lentiviral Particle

Symbol: Delta Opioid Receptor
Synonyms: DOP; DOR; DOR1; OPRD

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000911 **ORF Size:** 1116 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219275).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000911.3

 RefSeq Size:
 1774 bp

 RefSeq ORF:
 1119 bp

 Locus ID:
 4985

 UniProt ID:
 P41143

 Cytogenetics:
 1p35.3

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction





Delta Opioid Receptor (OPRD1) (NM_000911) Human Tagged ORF Clone Lentiviral Particle – RC219275L2V

MW: 40.2 kDa

Gene Summary:

G-protein coupled receptor that functions as receptor for endogenous enkephalins and for a subset of other opioids. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase. Signaling leads to the inhibition of adenylate cyclase activity. Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Plays a role in the perception of pain and in opiate-mediated analgesia. Plays a role in developing analgesic tolerance to morphine. [UniProtKB/Swiss-Prot Function]