

# Product datasheet for RC219347

## MRPL42 (NM\_172178) Human Tagged ORF Clone

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	MRPL42 (NM_172178) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	MRPL42
Synonyms:	HSPC204; MRP-L31; MRPL31; MRPS32; PTD007; RPML31
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>&gt;RC219347 representing NM_172178 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGGTGATGTCAAAGAGAACTATCTTGAAACATTTATTT
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	<pre>&gt;RC219347 representing NM_172178 Red=Cloning site Green=Tags(s)</pre>
	MGDVKENYLETFISSPKFGRTCADGALYCVCHKSTYSPLPDDYNCNVELALTSDGRTIVCYHPSVDIPYE HTKPIPRPDPVHNNEETHDQVLKTRLEEKVEHLEEGPMIEQLSKMFFTTKHRWYPHGRYHRCRKNLNPPK DR
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mk6323_f04.zip



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


## **GRIGENE** MRPL42 (NM\_172178) Human Tagged ORF Clone – RC219347

	Sgfl-Mlul
Cloning Scheme:	Cloning sites used for ORF Shuttling: Sgf I ORF Miu I GCGATCGC C ATG MINI ACG CGT
	Kozac         Consensus         EcoRI       BamHI Kpn I       RBS       SgfI         CTATAGGGCGGCCGGGAATTCGTCGGCTACCGGTACCGGGGAATCTGCCGCGCGCG
	$\frac{M[u] Noti Xhol }{T R T R P L R Q R L I S R E}$ $\frac{EcoR V}{Flag.Tag}$ Fileg.Tag $\frac{EcoR V}{T R T R P L R Q R L I S R E}$ $\frac{EcoR V}{T R T R R P L R Q R L I R R P R R$
	<u>DL</u> A A N D I L <u>D Y K D D D K</u> V Stop
CCN:	NM_172178
RF Size:	426 bp
TI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
TI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
omponents:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
econstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
efSeq:	<u>NM 172178.2, NP 751918.1</u>
	2093 bp
-	
efSeq Size: efSeq ORF:	428 bp


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	MRPL42 (NM_172178) Human Tagged ORF Clone – RC219347
Cytogenetics:	12q22
MW:	16.4 kDa
Gene Summary:	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a protein identified as belonging to both the 28S and the 39S subunits. Alternative splicing results in multiple transcript variants. Pseudogenes corresponding to this gene are found on chromosomes 4q, 6p, 6q, 7p, and 15q. [provided by RefSeq, May 2011]

## **Product images:**



Circular map for RC219347



Western blot validation of overexpression lysate (Cat# [LY406766]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC219347 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US