

## Product datasheet for RC219382L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PKM2 (PKM) (NM\_182470) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** PKM2 (PKM) (NM\_182470) Human Tagged ORF Clone Lentiviral Particle

Symbol: PKM2

Synonyms: CTHBP; HEL-S-30; OIP3; p58; PK3; PKM2; TCB; THBP1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM\_182470

ORF Size: 1593 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219382).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 182470.1

 RefSeq Size:
 2674 bp

 RefSeq ORF:
 1596 bp

 Locus ID:
 5315

 UniProt ID:
 P14618

 Cytogenetics:
 15q23

**Protein Families:** Druggable Genome





## PKM2 (PKM) (NM\_182470) Human Tagged ORF Clone Lentiviral Particle - RC219382L1V

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, Purine metabolism, Pyruvate metabolism,

Type II diabetes mellitus

**MW:** 57.9 kDa

**Gene Summary:** This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase

that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone

generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial

pathogenesis. Several alternatively spliced transcript variants encoding a few distinct

isoforms have been reported. [provided by RefSeq, May 2011]