

## Product datasheet for RC219394L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LATS2 (NM\_014572) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** LATS2 (NM\_014572) Human Tagged ORF Clone Lentiviral Particle

Symbol: LATS.
Synonyms: KPM

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_014572

 ORF Size:
 3264 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC219394).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 014572.1

 RefSeq Size:
 4098 bp

 RefSeq ORF:
 3267 bp

 Locus ID:
 26524

 UniProt ID:
 Q9NRM7

 Cytogenetics:
 13q12.11

**Domains:** UBA, pkinase, S\_TK\_X, TyrKc, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase





## LATS2 (NM\_014572) Human Tagged ORF Clone Lentiviral Particle - RC219394L1V

**MW:** 120 kDa

**Gene Summary:** 

This gene encodes a serine/threonine protein kinase belonging to the LATS tumor suppressor family. The protein localizes to centrosomes during interphase, and early and late metaphase. It interacts with the centrosomal proteins aurora-A and ajuba and is required for accumulation of gamma-tubulin and spindle formation at the onset of mitosis. It also interacts with a negative regulator of p53 and may function in a positive feedback loop with p53 that responds to cytoskeleton damage. Additionally, it can function as a co-repressor of androgen-responsive gene expression. [provided by RefSeq, Jul 2008]