

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC219403L2V

## INSIG1 (NM\_198336) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	INSIG1 (NM_198336) Human Tagged ORF Clone Lentiviral Particle
Symbol:	INSIG1
Synonyms:	CL6
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_198336
ORF Size:	990 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC219403).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 198336.1, NP 938150.1</u>
RefSeq Size:	3049 bp
RefSeq ORF:	537 bp
Locus ID:	3638
UniProt ID:	<u>015503</u>
Cytogenetics:	7q36.3
Protein Families:	Druggable Genome, Transmembrane
MW:	20.2 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes an endoplasmic reticulum membrane protein that regulates cholesterol<br/>metabolism, lipogenesis, and glucose homeostasis. The encoded protein has six<br/>transmembrane helices which contain an effector protein binding site. It binds the sterol-<br/>sensing domains of sterol regulatory element-binding protein (SREBP) cleavage-activating<br/>protein (SCAP) and 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG-CoA reductase),<br/>and is essential for the sterol-mediated trafficking of these two proteins. It promotes the<br/>endoplasmic reticulum retention of SCAP and the ubiquitin-mediated degradation of HMG-<br/>CoA reductase. Alternative splicing results in multiple transcript variants. [provided by<br/>RefSeq, Oct 2016]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US