

Product datasheet for RC219403L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

INSIG1 (NM_198336) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: INSIG1 (NM_198336) Human Tagged ORF Clone Lentiviral Particle

Symbol: INSIG1
Synonyms: CL6

Synonyms: CL6

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_198336

ORF Size: 990 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219403).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 198336.1, NP 938150.1

 RefSeq Size:
 3049 bp

 RefSeq ORF:
 537 bp

 Locus ID:
 3638

 UniProt ID:
 015503

Cytogenetics: 7q36.3

Protein Families: Druggable Genome, Transmembrane

MW: 20.2 kDa







Gene Summary:

This gene encodes an endoplasmic reticulum membrane protein that regulates cholesterol metabolism, lipogenesis, and glucose homeostasis. The encoded protein has six transmembrane helices which contain an effector protein binding site. It binds the sterolsensing domains of sterol regulatory element-binding protein (SREBP) cleavage-activating protein (SCAP) and 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG-CoA reductase), and is essential for the sterol-mediated trafficking of these two proteins. It promotes the endoplasmic reticulum retention of SCAP and the ubiquitin-mediated degradation of HMG-CoA reductase. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2016]