

Product datasheet for RC219408L4

CD40 (NM_152854) Human Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	CD40 (NM_152854) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	CD40
Synonyms:	Bp50; CDW40; p50; TNFRSF5
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC219408).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I ORF Mlu I GCG ATC GC ATG // NNN ACG CGT



ACCN: ORF Size: NM_152854 609 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CD40 (NM_152854) Human Tagged Lenti ORF Clone – RC219408L4	
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 152854.1</u>
RefSeq Size:	1115 bp
RefSeq ORF:	612 bp
Locus ID:	958
UniProt ID:	<u>P25942</u>
Cytogenetics:	20q13.12
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane
Protein Pathways:	Allograft rejection, Asthma, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Cytokine-cytokine receptor interaction, Primary immunodeficiency, Systemic lupus erythematosus, Toll-like receptor signaling pathway, Viral myocarditis
MW:	20.1 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD40 (NM_152854) Human Tagged Lenti ORF Clone – RC219408L4

Gene Summary:This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on
antigen-presenting cells of the immune system and is essential for mediating a broad variety
of immune and inflammatory responses including T cell-dependent immunoglobulin class
switching, memory B cell development, and germinal center formation. AT-hook transcription
factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand,
which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with
this receptor and serves as a mediator of the signal transduction. The interaction of this
receptor and its ligand is found to be necessary for amyloid-beta-induced microglial
activation, and thus is thought to be an early event in Alzheimer disease pathogenesis.
Mutations affecting this gene are the cause of autosomal recessive hyper-lgM
immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this
gene encoding distinct isoforms have been reported. [provided by RefSeq, Nov 2014]

Product images:



Circular map for RC219408L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US