

Product datasheet for RC219415L2

techsı EU: in

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

CD101 (NM_004258) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: CD101 (NM_004258) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: CD101

Synonyms: EWI-101; IGSF2; V7

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219415).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





 $[\]ensuremath{^*}$ The last codon before the Stop codon of the ORF.

ACCN: NM_004258

ORF Size: 3063 bp





OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

Domains:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

NM 004258.4 RefSeq:

RefSeq Size: 3557 bp RefSeq ORF: 3066 bp Locus ID: 9398 **UniProt ID:** Q93033 Cytogenetics: 1p13.1

Protein Families: Transmembrane

ig, IGv, IG

MW: 115.1 kDa

Plays a role as inhibitor of T-cells proliferation induced by CD3. Inhibits expression of IL2RA **Gene Summary:**

on activated T-cells and secretion of IL2. Inhibits tyrosine kinases that are required for IL2


production and cellular proliferation. Inhibits phospholipase C-gamma-1/PLCG1

phosphorylation and subsequent CD3-induced changes in intracellular free calcium. Prevents nuclear translocation of nuclear factor of activated T-cell to the nucleus. Plays a role in the inhibition of T-cell proliferation via IL10 secretion by cutaneous dendritic cells. May be a

marker of CD4(+) CD56(+) leukemic tumor cells.[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for RC219415L2