

Product datasheet for RC219430L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LCK (NM_001042771) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LCK (NM_001042771) Human Tagged ORF Clone Lentiviral Particle

Symbol: LCK

Synonyms: IMD22; LSK; p56lck; pp58lck; YT16

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001042771

ORF Size: 1527 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219430).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001042771.1, NP 001036236.1

 RefSeq Size:
 2118 bp

 RefSeq ORF:
 1530 bp

 Locus ID:
 3932

 UniProt ID:
 P06239

Cytogenetics: 1p35.2

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency



LCK (NM_001042771) Human Tagged ORF Clone Lentiviral Particle - RC219430L4V

Protein Pathways: Natural killer cell mediated cytotoxicity, Primary immunodeficiency, T cell receptor signaling

pathway

MW: 57.8 kDa

Gene Summary: This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded

protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-

containing and proline-rich motifs, respectively. The protein localizes to the plasma

membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants encoding

different isoforms have been described. [provided by RefSeq, Aug 2016]