

Product datasheet for RC219509L3

FNIP2 (NM_020840) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	FNIP2 (NM_020840) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	FNIP2
Synonyms:	FNIP1; MAPO1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC219509).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_020840
ORF Size:	3342 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_020840.1
RefSeq Size:	5801 bp
RefSeq ORF:	3345 bp
Locus ID:	57600
UniProt ID:	Q9P278
Cytogenetics:	4q32.1
MW:	121.9 kDa
Gene Summary:	This gene encodes a protein that binds to the tumor suppressor folliculin and to AMP-activated protein kinase (AMPK), and may play a role cellular metabolism and nutrient sensing by regulating the AMPK-mechanistic target of rapamycin signaling pathway. The encoded protein may also be involved in regulating the O6-methylguanine-induced apoptosis signaling pathway. This gene has a closely related paralog that encodes a protein with similar binding activities. Both related proteins also associate with the molecular chaperone heat shock protein-90 (Hsp90) and negatively regulate its ATPase activity and facilitate its association with folliculin. [provided by RefSeq, Jul 2017]

Product images:



Circular map for RC219509L3