

Product datasheet for RC219579L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NIRF (UHRF2) (NM_152896) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NIRF (UHRF2) (NM 152896) Human Tagged ORF Clone Lentiviral Particle

Symbol: NIRF

Synonyms: NIRF; RNF107; TDRD23; URF2

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 152896

Tag: Myc-DDK

ORF Size: 2406 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219579).

•

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. <u>More info</u>

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 152896.1

 RefSeq Size:
 3621 bp

 RefSeq ORF:
 2409 bp

 Locus ID:
 115426

 UniProt ID:
 Q96PU4

 Cytogenetics:
 9p24.1

Protein Families: Druggable Genome, Transcription Factors

MW: 89.8 kDa







Gene Summary:

This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. The encoded protein contains an NIRF_N domain, a PHD finger, a set- and ring-associated (SRA) domain, and a RING finger domain and several of these domains have been shown to be essential for the regulation of cell proliferation. This protein may also have a role in intranuclear degradation of polyglutamine aggregates. Alternative splicing results in multiple transcript variants some of which are non-protein coding. [provided by RefSeq, Feb 2012]