

## Product datasheet for RC219628L4V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Vitamin D Receptor (VDR) (NM 001017535) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Vitamin D Receptor (VDR) (NM\_001017535) Human Tagged ORF Clone Lentiviral Particle

Symbol: Vitamin D Receptor Synonyms: NR1I1; PPP1R163

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001017535

ORF Size: 1281 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219628).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001017535.1

 RefSeq Size:
 4791 bp

 RefSeq ORF:
 1284 bp

 Locus ID:
 7421

 UniProt ID:
 P11473

 Cytogenetics:
 12q13.11

**Protein Families:** Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

MW: 48.1 kDa





# Vitamin D Receptor (VDR) (NM\_001017535) Human Tagged ORF Clone Lentiviral Particle – RC219628L4V

#### **Gene Summary:**

This gene encodes vitamin D3 receptor, which is a member of the nuclear hormone receptor superfamily of ligand-inducible transcription factors. This receptor also functions as a receptor for the secondary bile acid, lithocholic acid. Downstream targets of vitamin D3 receptor are principally involved in mineral metabolism, though this receptor regulates a variety of other metabolic pathways, such as those involved in immune response and cancer. Mutations in this gene are associated with type II vitamin D-resistant rickets. A single nucleotide polymorphism in the initiation codon results in an alternate translation start site three codons downstream. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon. [provided by RefSeq, Jun 2018]