

Product datasheet for RC219641L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SCDGFB (PDGFD) (NM_025208) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SCDGFB (PDGFD) (NM_025208) Human Tagged ORF Clone Lentiviral Particle

Symbol: SCDGFB

Synonyms: IEGF; MSTP036; SCDGF-B; SCDGFB

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 025208

ORF Size: 1110 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC219641).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 025208.3

 RefSeq Size:
 3870 bp

 RefSeq ORF:
 1113 bp

 Locus ID:
 80310

 UniProt ID:
 Q9GZP0

 Cytogenetics:
 11q22.3

Domains: CUB

Protein Families: Druggable Genome, Secreted Protein





SCDGFB (PDGFD) (NM_025208) Human Tagged ORF Clone Lentiviral Particle - RC219641L3V

Protein Pathways: Focal adhesion, Gap junction, Melanoma, Prostate cancer, Regulation of actin cytoskeleton

MW: 40.2 kDa

Gene Summary: The protein encoded by this gene is a member of the platelet-derived growth factor family.

The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a core motif of eight cysteines, seven of which are found in this factor. This gene product only forms homodimers and, therefore, does not dimerize with the other three family members. It differs from alpha and beta members of this family in having an unusual N-terminal domain, the CUB domain. Two splice variants have been identified for

this gene. [provided by RefSeq, Jul 2008]