

## Product datasheet for RC219686L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **TEAD4 (NM\_003213) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** TEAD4 (NM\_003213) Human Tagged ORF Clone Lentiviral Particle

Symbol: TEAD4

Synonyms: EFTR-2; hRTEF-1B; RTEF1; TCF13L1; TEF-3; TEF3; TEFR-1

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_003213

ORF Size: 1302 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219686).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 003213.2

 RefSeq Size:
 1798 bp

 RefSeq ORF:
 1305 bp

 Locus ID:
 7004

 UniProt ID:
 Q15561

Cytogenetics: 12p13.33

Domains: TEA

**Protein Families:** Transcription Factors





ORIGENE

MW: 48.3 kDa

**Gene Summary:** This gene product is a member of the transcriptional enhancer factor (TEF) family of

transcription factors, which contain the TEA/ATTS DNA-binding domain. It is preferentially expressed in the skeletal muscle, and binds to the M-CAT regulatory element found in promoters of muscle-specific genes to direct their gene expression. Alternatively spliced transcripts encoding distinct isoforms, some of which are translated through the use of a non-AUG (UUG) initiation codon, have been described for this gene. [provided by RefSeq, Jul

2008]