

Product datasheet for RC219839L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FAK (PTK2) (NM_153831) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: FAK (PTK2) (NM 153831) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: FAK

Synonyms: FADK; FADK 1; FAK; FAK1; FRNK; p125FAK; pp125FAK; PPP1R71

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC219839).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





st The last codon before the Stop codon of the ORF.

ACCN: NM_153831

ORF Size: 3165 bp





FAK (PTK2) (NM_153831) Human Tagged Lenti ORF Clone - RC219839L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 153831.2</u>

 RefSeq Size:
 4453 bp

 RefSeq ORF:
 3159 bp

 Locus ID:
 5747

UniProt ID: Q05397

Cytogenetics: 8q24.3

Domains:B41, pkinase, TyrKc, S_TKc, Focal_AT **Protein Families:**Druggable Genome, Protein Kinase

Protein Pathways: Axon guidance, Chemokine signaling pathway, ErbB signaling pathway, Focal adhesion,

Leukocyte transendothelial migration, Pathways in cancer, Regulation of actin cytoskeleton,

Small cell lung cancer, VEGF signaling pathway

MW: 119.1 kDa

Gene Summary: This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the

focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation


of this gene may be an important early step in cell growth and intracellular signal

transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have

been found for this gene. [provided by RefSeq, Jun 2017]



Product images:



Double digestion of RC219839L3 using Sgfl and Mlul $\,$