

## Product datasheet for **RC219845L1V**

### ACCN1 (ASIC2) (NM\_183377) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	ACCN1 (ASIC2) (NM_183377) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ACCN1
Synonyms:	ACCN; ACCN1; ASIC2a; BNaC1; BNC1; hBNaC1; MDEG
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_183377
ORF Size:	1689 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC219845).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_183377.1</a> , <a href="#">NP_899233.1</a>
RefSeq Size:	3470 bp
RefSeq ORF:	1692 bp
Locus ID:	40
UniProt ID:	<a href="#">Q16515</a>
Cytogenetics:	17q11.2-q12
Protein Families:	Druggable Genome, Ion Channels: Other
Protein Pathways:	Taste transduction



[View online »](#)

**MW:** 62.7 kDa

**Gene Summary:** This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, 2 hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene may play a role in neurotransmission. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 3 has been observed to co-assemble into proton-gated channels sensitive to gadolinium. Alternative splicing has been observed at this locus and two variants, encoding distinct isoforms, have been identified. [provided by RefSeq, Feb 2012]