

## Product datasheet for RC219892L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PIGC (NM\_002642) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** PIGC (NM\_002642) Human Tagged ORF Clone Lentiviral Particle

Symbol: PIGC

**Synonyms:** GPI2; GPIBD16; MRT62

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 002642

ORF Size: 891 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC219892).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002642.3

 RefSeq Size:
 1537 bp

 RefSeq ORF:
 894 bp

 Locus ID:
 5279

 UniProt ID:
 Q92535

 Cytogenetics:
 1q24.3

**Protein Families:** Transmembrane

**Protein Pathways:** Glycosylphosphatidylinositol(GPI)-anchor biosynthesis, Metabolic pathways





**MW:** 33.6 kDa

**Gene Summary:** 

This gene encodes an endoplasmic reticulum associated protein that is involved in glycosylphosphatidylinositol (GPI) lipid anchor biosynthesis. The GPI lipid anchor is a glycolipid found on many blood cells and serves to anchor proteins to the cell surface. The encoded protein is one subunit of the GPI N-acetylglucosaminyl (GlcNAc) transferase that transfers GlcNAc to phosphatidylinositol (PI) on the cytoplasmic side of the endoplasmic reticulum. Two alternatively spliced transcripts that encode the same protein have been found for this gene. A pseudogene on chromosome 11 has also been characterized. [provided by RefSeq, Jul 2008]