

Product datasheet for **RC219931**

DCAMKL2 (DCLK2) (NM_001040260) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	DCAMKL2 (DCLK2) (NM_001040260) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	DCAMKL2
Synonyms:	CL2; CLICK-II; CLICK2; CLIK2; DCAMKL2; DCDC3; DCDC3B; DCK2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RC219931 representing NM_001040260
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCCAGCACCAGGAGTATCGAGCTGGAGCACTTTGAGGAACGGGACAAAAGGCCGGCCGGGGTCCG
 GGAGAGGGGCCCCAGCTCCTCCGGGGCAGCAGCAGCTCGGGCCCAAGGGGAACGGGCTCATCCCCAG
 TCCGGCGCACAGTGCCCACTGCAGCTTCTACCGCACGCGACCCTGCAGGCCCTCAGCTCGGAGAAGAAG
 GCCAAGAAGGCGCGCTTCTACCGGAACGGGGACCCTACTTCAAGGGCCTGGTGTGGCCATCTCCAGCG
 ACCGCTCCGGTCTTCGATGCGCTCCTCATAGAGCTCACCCGCTCCCTGTCGGACAACGTGAACCTGCC
 CCAGGGTGTCCGCACTATCTACACCATCGACGGCAGCCGGAAGGTACCAGCCTGGACGAGCTGCTGGAA
 GGTGAGAGTTACGTGTGCATCCAATGAACATTTTCGTAAGTCGATTACACAAAAATTAATCCAA
 ACTGGTCTGTGAACATCAAGGGTGGGACATCCCGAGCGCTGGCTGCTGCCTCCTCTGTGAAAAGTGAAGT
 AAAAGAAAGTAAAGATTTTCATCAAAACCAAGTTAGTGACTGTGATTTCGAAGTGGAGTGAAGCCTAGAAA
 GCCGTGCCGATCCTTCTGAATAAAAAGACTGCTCATTCTTTGAACAAGTCTTAACAGATATCACCGAAG
 CCATTAATACTAGACTCAGGAGTCGTAAGAGGCTCTGCACCCTGGATGGAAGCAGGTTACTTGTCTGCA
 AGACTTTTTGGTGATGACGATGTTTTATTGCATGTGGACCAGAAAAATTCGTTATGCCCAAGATGAC
 TTTGTCCTGGATCATAGTGAATGTCGTGCTGAAGTCATCTTATTCTCGATCCTCAGCTGTTAAGTATT
 CTGGATCCAAAAGCCCTGGGCCCTCTCGACGCAGCAATCACCAGCTTCAGTTAATGGAACCTCCAGCAG
 CCAACTTTCTACTCCTAAATCTACGAAATCCTCCAGTTCCTCTCCAAGTCCAGGAAGTTTCAGAGGA
 TAAAGCAGATTTCTGCTCATGGCAGATCTTCTCCAATGTAAACGGTGGACCTGAGCTTGACCAATTGCA
 TAAGCTGAAGGTGTGAATGGAACAGATGCTCTGAATCATCAACTCTTCTTGAGAAATACAAAATTGG
 AAAGGTCAATGGTGATGGCAATTTTGCAGTAGTCAAAGAGTGTATAGACAGGTCCACTGGAAAGGATTT
 GCCCTAAAGATTATAGACAAAGCCAAATGTTGTGAAAGGAACACCTGATTGAGAATGAAGTGTCAATAC
 TGCGCCGAGTGAAACATCCCAATATCATTATGCTGGTTCGAGGAGATGGAACAGCAACTGAGCTCTTTCT
 GGTGATGGAATTGGTCAAAGGTGGAGATCTCTTTGATGCAATTACTTCGTCGACCAAGTACACTGAGAGA
 GATGGCAGTGCCATGGTGTACAACCTAGCCAATGCCCTCAGGTATCTCCATGGCCTCAGCATCGTGACA
 GAGACATCAAACCAGAGAATCTTGGTGTGTAATATCCTGATGGAACCAAGTCTTTGAAACTGGGAGA
 CTTTGGGCTTGCAGCTGTGGTAGAAGGCCCTTTATACACAGTCTGTGGCACACCCACTTATGTGGCTCCA
 GAAATCATTGCTGAAACTGGCTATGGCCTGAAGGTGGACATTTGGGCAGCTGGTGTGATCACATACATAC
 TTCTCTGTGGATCCCACCATCCGAAGTGAGAACAATCTCCAGGAAGATCTCTTCGACCAGATCTTGGC
 TGGGAAGCTGGAGTTCCGGCCCCCTACTGGGATAACATCACGGACTCTGCCAAGGAATTAATCAGTCAA
 ATGCTTACAGTAAATGTTGAAGCTCGGTGTACCGCGGGACAAATCCTGAGTCAACCCTGGGTGTCAGATG
 ATGCCCTCCAGGAGAATAACATGCAAGCTGAGGTGACAGGTAAACTAAAACAGCACTTTAATAATGCGCT
 CCCCAACAGAACAGCACTACCACCGGGTCTCCGTCATCATGAACACGGCTCTAGATAAGGAGGGGCGAG
 ATTTTCTGCAGCAAGCACTGTCAAGACAGCGGCAGGCCTGGGATGGAGCCCATCTCTCCAGTTCCCTCCT
 CAGTGGAGGAGATCCCTGTGCTGGGGAAGCAGTCCCGGCCCCACCCCTCCGGAATCTCCCACCCCCCA
 CTGTCTCCCGCTGCCCCGGTGGTGTGAGCGGGCAGGAACCTGGCGCCGCCACCGAGAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC219931 representing NM_001040260
Red=Cloning site Green=Tags(s)

MASTRSIELEHFEERDKRPRPGSRRGAPSSSSGGSSSSGPKGNGLIPSPAHSACSFYRTRTLQALSSEKK
 AKKARFYRNGDRYFKGLVFAISSDRFRSFDALLIELTRSLSDNVNLPQGVRTIYIDGSRKVTSLDELLE
 GESYVCASNEPFRKVDYTKNINPNWSVNIKGGTSRALAAASSVKSEVKESKDFIKPKLVTIRSGVKPRK
 AVRILLNKTAHSFEQVLTDITEAIKLD SGVVKRLCTLDGKQVTC LQDFFGDDDFVIACGPEKFRYAQDD
 FVLDHSECRVLKSSYSRSSAVKYSGSKSPGSRRSKSPASVNGTPSSQLSTPKSTKSSSSSPTSPGSFRG
 LKQISAHGRSSSNVNGGPELDRCSPEGVNGNRCSESSTLLEKYKIGKVI GDGNFAVVKECIDRSTGKEF
 ALKIIDKAKCCGKEHLIENEVSILRRVKHPNIIMLVEEMETATELFLVMELVKGGDLFDAITSSTKYTER
 DGSAMVYNLANALRYLHGLSIVHRDIKPENLLVCEYPDGTSKSLKLGDFLATVVEGPLYTVCGTPTYVAP
 EIIAETGYLKVDIWAAGVITYILLCGFPPFRSENQLQEDLFDQILAGKLEFPAPYWDNITDSAKELISQ
 MLQVNVEARCTAGQILSHPWVSDASQENNMQAEVTGKLGKQHFNNALPKQNSTTTGVSVIMNTALDKEGQ
 IFCSKHCQDSGRPGMEIIPVPPSVEEIPVPGEA VPA TPPE SPTPHCPPAAPGGERAGTWRRHRD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_001040260

ORF Size: 2298 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001040260.1](#), [NP_001035350.1](#)

RefSeq Size: 3603 bp

RefSeq ORF: 2301 bp

Locus ID: 166614

UniProt ID: [Q8N568](#)


Cytogenetics: 4q31.23-q31.3

Protein Families: Druggable Genome, Protein Kinase


MW: 83.4 kDa

Gene Summary: This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca²⁺/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family member, and this gene share function in the establishment of hippocampal organization and that their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alternatively spliced transcript variants have been identified. [provided by RefSeq, Sep 2010]

Product images:



Circular map for RC219931



Western blot validation of overexpression lysate (Cat# [LY421729]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC219931 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

□

Coomassie blue staining of purified DCLK2 protein (Cat# [TP319931]). The protein was produced from HEK293T cells transfected with DCLK2 cDNA clone (Cat# RC219931) using MegaTran 2.0 (Cat# [TT210002]).